

Communities of Microorganisms in Their Natural Setting

Hong An Dao
Department of Biotechnology
HO CHI MINH CITY OPEN UNIVERSITY
Ho Chi Minh City, Vietnam

Abstract

Aeromicrobiology investigates microorganisms carried through the air, highlighting their lower quantities than seas and soil. However, once dispersed, these infectious agents can travel long distances, increasing the risk of epidemics and impacting the environment and health.

Keywords: *microorganisms, airborne, epidemics, environment, health, bacteria, fungi, cloud chemistry, bioaerosols*

Introduction

Aeromicrobiology, the investigation of microorganisms carried through the air, highlights that airborne microorganism quantities are significantly lower than those found in seas and soil. However, once dispersed in the atmosphere, these infectious agents can travel long distances aided by wind and storms, significantly raising the risk of epidemics. These aerosols have environmental impacts and are linked to various health problems affecting all life forms.

Bacteria, frequently observed floating in clouds, can modify cloud chemistry through their metabolic processes, potentially leading to precipitation formation. Multiple physical processes influence the release, movement, and collection of bioaerosols. Airborne particles can originate from land or water habitats, with turbulence propelling them higher into the sky. Wind plays a crucial role in transporting bioaerosols between locations, while gravity, direct contact with surfaces, or interactions with precipitation bring them back to the Earth's surface.

Germes and other particles like water droplets, dirt, and debris are airborne. Bacteria follow a specific path and can be propelled into the air. Microbes in the air stem from various sources, including humans, animals, and plants, and are transported to new environments through different vectors like wind, equipment, and people. Atmospheric conditions, such as temperature, relative humidity, and radiation levels, significantly impact the types of microorganisms that can survive and the duration of their presence in the air.

Physical surroundings strains

Atmospheric conditions constrain bacteria, the greatest being the risk of desiccation when in free fall. Their survival is also affected by the humidity in the air, with Gram-negative microorganisms being more tolerant of damp conditions than their Gram-positive counterparts. Proteins may get denatured, or ice crystals can develop at frigid temperatures. Radiation is also dangerous since it has been shown to cause DNA damage in aeromicrobes.

Microbial communities

Atmospheric microbial communities are made up of many different kinds of microorganisms that exist as aerosols in the air, such as viruses, bacteria, fungi, yeasts, and protozoans. These bacteria, to survive, must adjust to challenging environments, including high humidity, high particulate matter, and low moisture. Many bacteria that can live in such settings also have the ability to create spores that can survive in even harsher environments.

Antimicrobial Air Pollution refers to microorganisms in the air that harm humans and other species. These microorganisms include viruses, bacteria, and fungus. Due to the turbulence in the atmosphere, these microorganisms are dispersed from places like disturbed soil, plants, and ocean surfaces. Different variables, such as time of day, season, location, and sources, affect the concentration of microorganisms in the air. They may scatter rapidly or float for great distances. Many airborne cells may be rendered nonviable by variations in temperature, hydration, recovery, exposure to UV light, and air pollution. These impacts may be simulated using mathematical models that also examine microbial survival.

Bacterial Species Surviving in Harsh Environments:

Bacillus anthracis, a gram-positive, rod-shaped bacterium, has the remarkable ability to thrive in harsh conditions. It defends itself by producing spores, which are dormant structures with thick cell membranes that help withstand adverse environmental factors. *Bacillus anthracis* can withstand high temperatures, chemical pollution, and nutrient scarcity, making it a resilient bacterium. Notably, this bacterial species is associated with anthrax, a deadly form of pneumonia.

Fungal Pathogen Adapted to Harsh Environments:

Aspergillus fumigatus, a common airborne fungal pathogen, is another example of a microbe capable of thriving in challenging environments. When inhaled, its spores or conidia can cause various illnesses in humans. Despite lacking specific pathogenic traits, *Aspergillus fumigatus* can cause widespread infections due to its impressive adaptability to diverse environmental conditions.

The Challenges of the Environment:

The environment presents significant challenges due to its highly psychrophilic and concentrated nature. Cloud temperatures average below 0°C, and atmospheric humidity has been recorded as low as -101 degrees Celsius. Despite these extreme conditions, scientists have discovered many atmospheric microbes, including over 103 bacterial species. Remarkably, microorganisms capable of growth and multiplication have been isolated from cloud water, even at altitudes as high as 78 km in the stratosphere.

Expanding Understanding of Microbial Roles:

Research is expanding our understanding of microbes' roles in various processes, such as ice formation, cloud formation, ammonia metabolic processes, and the decomposition of carbon-based organic compounds. However, the lack of consensus on optimal bioaerosol sampling procedures has impeded progress in this field. Although aerobiological studies have been conducted since the eighteenth century, our comprehension of atmospheric microbes is still in its early stages. As we recognize the importance of understanding microbial biogeography, further research is needed to explore the practical implications of psychrophilic microbes in the environment. This research is relevant in biological geography, long-range meteorological propagation, human health, and agriculture.

Conclusion

Aeromicrobiology emphasizes the lower quantities of airborne microorganisms but recognizes their ability to travel long distances, posing risks to epidemics and environmental health. Bacteria and fungi exhibit adaptability to harsh environments. Research on atmospheric microbes and their roles in processes like precipitation formation is advancing, yet the lack of consensus on sampling procedures hinders progress. Further exploration of psychrophilic microbes is crucial for understanding physical geography, meteorological propagation, human health, and agriculture.

References

1. <https://microbewiki.kenyon.edu/index.php/Aeromicrobiology>
2. Tina Šantl-Temkiv and others, Microbial ecology of the atmosphere, *FEMS Microbiology Reviews*, Volume 46, Issue 4, July 2022, fuac009, <https://doi.org/10.1093/femsre/fuac009>
3. Chen X, Kumari D, Achal V. A Review on Airborne Microbes: The Characteristics of Sources, Pathogenicity and Geography. *Atmosphere*. 2020; 11(9):919. <https://doi.org/10.3390/atmos11090919>